

Faculty of Engineering & Technology

First Semester B.E. (C.B.S.) Examination

APPLIED MATHEMATICS—I

Paper—I

Time : Three Hours]

[Maximum Marks : 80]

INSTRUCTIONS TO CANDIDATES

- (1) All questions carry marks as indicated.
- (2) Use of non-programmable calculator is permitted.
- (3) Solve :

Question No. 1 OR Question No. 2

Question No. 3 OR Question No. 4

Question No. 5 OR Question No. 6

Question No. 7 OR Question No. 8

Question No. 9 OR Question No. 10

Question No. 11 OR Question No. 12.

1. (a) If $y = a \cos(\log x) + b \sin(\log x)$ show that :

$$x^2 y_{n+2} + (2n+1)xy_{n+1} + (n^2 + 1)y_n = 0. \quad 6$$

(b) Evaluate : $\lim_{x \rightarrow 0} \frac{x \cos x - \sin x}{x^2 \sin x}. \quad \text{Ans: } 0 \quad 3.$

(c) Evaluate : $\lim_{x \rightarrow 0} \left(\frac{a^x + b^x + c^x}{3} \right)^{1/x}. \quad \text{Ans: } (abc)^{1/3}$

OR $\log(abc)^{1/3}$

2. (a) If $x = a \cos^4 \theta, y = a \sin^4 \theta$; find the curvature at $\theta = \pi/6. \quad 6$

- (b) Using Taylor's series find the value of $\cos 64^\circ$ correct to four decimal places. 6

3. (a) If $u(x+y) = x^2 + y^2,$

then prove that :

$$\left(\frac{\partial u}{\partial x} - \frac{\partial u}{\partial y} \right)^2 = 4 \left(1 - \frac{\partial u}{\partial x} - \frac{\partial u}{\partial y} \right). \quad 6$$

- (b) If $u = \tan^{-1} \left[\frac{x^3 + y^3}{x-y} \right]$ prove that :

$$x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = \sin 4u - \sin 2u. \quad 6$$

- (c) If $u = f(x/y, y/z, z/x)$ find the value of :

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z}. \quad 4 \quad 2$$

OR

4. (a) Given $u = \frac{x-y}{x+y}, v = \frac{x+y}{x}$,

find $\frac{\partial(u,v)}{\partial(x,y)}$. Are u and v functionally related ?

If so, find the relation between them. 6

- (b) Expand y^x in the neighbourhood of $(1, 1)$ upto the term of second degree. 6

- (c) Find the points on the surface $z^2 = xy + 1$ nearest to origin. 6

5. (a) Find the inverse of matrix by partitioning method :

$$A = \begin{bmatrix} 2 & 3 & 4 \\ 4 & 3 & 1 \\ 1 & 2 & 4 \end{bmatrix}. \quad 6$$

Grat + 2+4+2- 211 x 21

(b) Find the rank of the matrix :

$$6 \quad A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 2 & 1 & 4 & 5 \\ 1 & 5 & 5 & 7 \\ 8 & 1 & 14 & 17 \end{bmatrix} \quad 6$$

OR

6. (a) Test the consistency and solve :

$$\begin{aligned} x + y + z &= 3 & -8x + 12z &= 6 \\ x + 2y + 3z &= 4 \\ x + 4y + 9z &= 6. \end{aligned}$$

(b) Solve the system of equations by adjoint method :

$$\begin{aligned} x - 2y + 3z &= 2 & 211615 \\ 2x - 3z &= 3 \\ x + y + z &= 0. \end{aligned}$$

7. (a) Solve : $(x+1) \frac{dy}{dx} - 2y = (x+1)^4$. 4

(b) Solve : $(1+x) \frac{dy}{dx} - \tan y = (1+x)^2 e^x \sec y$. 4

(c) Solve : $\frac{dy}{dx} + \frac{x+y \cos x}{1+\sin x} = 0$. 4

OR

8. (a) Solve : $P(P+y) = x(x+y)$. 4

(b) Solve : $y = 2px + p^n$, where $P = \frac{dy}{dx}$. 4

(c) Solve : $P^3 - 4xyP + 8y^2 = 0$. 4

9. (a) Solve : $\frac{d^2y}{dx^2} + 4y = \cos 2x + e^{3x}$. 6

(b) Solve by method of variation of parameter :

$$\frac{d^2y}{dx^2} - 6 \frac{dy}{dx} + 9y = \frac{e^{3x}}{x^2}. \quad 6$$

(c) Solve :

$$x^2 \frac{d^2y}{dx^2} - 3x \frac{dy}{dx} + 5y = x \log x. \quad 6$$

OR

10. (a) Solve :

$$\frac{dx}{dt} + 2x - 3y = t$$

$$\frac{dy}{dt} - 3x + 2y = e^{2t}. \quad 6$$

(b) Solve :

$$\frac{d^2y}{dx^2} = 3\sqrt{y}, \text{ given that}$$

$$y = 1, \frac{dy}{dx} = 2, \text{ when } x = 0. \quad 6$$

- (c) In an L-C-R circuit, the charge q on a plate of a condenser is given by $L \frac{d^2q}{dt^2} + R \frac{dq}{dt} + \frac{q}{C} = E \sin pt.$ The circuit is tuned to resonance so that $P^2 = \frac{1}{LC}.$ If initially current i and the charge q be zero, show that for small values of $R/L,$ the current in the circuit at time t is given by $\left(\frac{Et}{2L}\right) \sin pt.$ 6

11. (a) If $\tan(\theta + i\phi) = \cos \alpha + i \sin \alpha,$ prove that :

$$\theta = \frac{n\pi}{2} + \frac{\pi}{4} \text{ and } \phi = \frac{1}{2} \log \tan \left(\frac{\pi + \alpha}{4} \right). \quad 4$$

- (b) Find all the values of $\left(\frac{1}{2} + \frac{\sqrt{3}i}{2}\right)^{3/4}$ and show that their continual product is 1. 4

OR

6

(Contd.)

12. (a) Using De-Moivre's theorem, solve :

$$x^5 + x^4 + x^3 + x^2 + x + 1 = 0. \quad 4$$

- (b) If $2 \cos \theta = x + \frac{1}{x}, 2 \cos \phi = y + \frac{1}{y},$ show that :

$$x^m y^n + \frac{1}{x^m y^n} = 2 \cos(m\theta + n\phi). \quad 4$$