

Engineering Physics

P. Pages : 2

NKT/KS/17/7197

Time : Two Hours

* 0 2 3 1 *

Max. Marks : 40

- Notes :
1. All questions carry marks as indicated.
 2. Solve Question 1 OR Questions No. 2.
 3. Solve Question 3 OR Questions No. 4.
 4. Solve Question 5 OR Questions No. 6.
 5. Solve Question 7 OR Questions No. 8.
 6. Assume suitable data whenever necessary.
 7. Use of non programmable calculator is permitted.

List of constants :

1. Planck's constant ' h ' = 6.63×10^{-34} JS
2. Velocity of light ' C ' = 3×10^8 m/s
3. Charge on electron ' e ' = 1.602×10^{-19} C
4. Mass of electron ' m ' = 9.11×10^{-31} kg
5. Avogadro's constant ' N_A ' = $6.023 \times 10^{26} \frac{\text{atoms}}{\text{kmole}}$
6. Boltzman's constant ' K ' = 1.380×10^{-23} JK⁻¹

1. a) What is Compton effect? Why classical theory failed to explain it? 2+2
- b) In case of high atomic no. scatterer element intensity of unmodified wavelength is higher than that of modified wavelength. Explain. 3
- c) X-rays of initial wavelength 0.5×10^{-10} m undergo Compton Scattering. Find the scattering angle at which wavelength of scattered X-rays will be greater than that of incident wavelength by one percent? 3

OR

2. a) What is de-Broglie hypothesis? Obtain an expression for wavelength associated with an electron accelerated through a potential difference of ' V ' volts. 4
- b) Show how the Bohr's quantization condition of angular momentum follows the concept of matter waves. 3
- c) Calculate de-Broglie wavelength of the orbital electron of Hydrogen atom. (Given that energy of electron is 13.6 eV) 3
3. a) Derive an expression for wave function of an electron confined to move in an infinite potential well of width ' L '. 5
- b) What is a wave packet? 2

- c) Find the lowest energy of an electron in one dimensional potential well of width 2 \AA . Express the result in electron volts. **3**
4. a) Discuss a thought experiment of electron diffraction to arrive at Heisenberg uncertainty principle with suitable diagram. **4**
- b) Define phase velocity and group velocity. **2**
- c) Calculate uncertainty in location of an electron and a ball of mass 1 kg if their velocities are 10^5 m/s and 10 m/s respectively. **4**
5. a) Derive expression for atomic radius and atomic packing fraction (A.P.F.) for BCC and FCC structures and show that percentage void space is more in BCC compared to FCC structure. **6**
- b) Molybdenum has BCC structure. Its density is $10.2 \times 10^3 \text{ kg/m}^3$ and its atomic wt. is 95.94. Determine radius of Molybdenum atom. **4**
6. a) What are Miller Indices? Draw the crystal planes in simple cubic structure having Miller Indices (i) (102) and (ii) (0 10). **3**
- b) State and derive Bragg's Law of X-ray diffraction. **4**
- c) The d_{110} interplaner spacing in a BCC metal vanadium is 2.15 \AA . Find its lattice constant (a). **3**
7. a) Explain classification of solids on the basis of energy band diagrams. **3**
- b) Draw energy band diagrams for the following : **2+2**
- i) PN Junction in Reverse Bias
- ii) NPN Transistor (unbiased)
- c) Find V_0 across a silicon junction at room temperature, if p-region has 10^{21} acceptor atoms/ m^3 and N-region has 10^{22} atoms/ m^3 . **3**
8. a) What is Hall effect? Obtain an expression for Hall voltage if p-type semiconductor material is used. **2+3**
- b) What is meant by depletion region? **2**
- c) Determine the probability of an electron thermally excited into the conduction band in Germanium at 27°C , if the energy gap is 0.72 eV . **3**
